

Dietary Boron: Progress in Establishing Essential Roles in Human and Animal Physiology

C. Hunt

U.S. Dept. Agriculture, Agricultural Research Service, Grand Forks Human Nutrition Research Center, Grand Forks, ND, USA

ABSTRACT: This review summarizes the progress made in establishing essential roles for boron in human and animal physiology and assesses that progress in view of criteria for essentiality of elements. The evidence to date suggests that humans and at least some higher animals may use boron to support normal biological functions. These include roles in calcium metabolism, bone growth and maintenance, insulin metabolism, and completion of the life cycle. The biochemical mechanisms responsible for these effects are poorly understood. However, the nature of boron biochemistry suggests specific lines of investigation. In particular, further characterization of the cell signalling molecules capable of complexing with boron should provide insights into the specific biochemical function(s) of boron in humans.

1 INTRODUCTION


Boron has long been recognized as essential for all higher plants in phylogenetic kingdom Viridiplantae (Warington, 1926). More recently, boron was recognized as required for at least some organisms in Eubacteria (Chen et al., 1981), Stramenopila (Lovatt, 1985), and Animalia (Fort et al., 2002; Rowe, 1999). Specific species in the Fungi kingdom have a demonstrated physiological response to boron, an important finding because Fungi species are thought to share a common ancestor with animals exclusive of plants (Carney et al., 2004). This review summarizes the progress made in establishing essential roles for boron in human and animal physiology and assesses that progress in relation to accepted criteria for essentiality of elements (Expert Consultation WHO/FAO/IAEA, 1996; Frieden, 1984; Mertz, 1970; Underwood et al., 1987). Those criteria are: 1) it reacts with biological material or forms chelates; 2) it is present in healthy tissues of different animals at comparable concentrations; 3) toxicity results only at relatively high intakes; 4) tissue concentrations during short term variations in intake are maintained by homeostatic mechanisms; 5) depletion prevents growth and completion of the life cycle; 6) depletion consistently results in reduction of a physiologically important function; and 7) when an integral part of an organic structure, depletion causes reduction in performance of a vital function.

2 BORON BIOMOLECULES

2.1 Boron chemistry

Organoboron compounds are those organic compounds that contain B-O bonds, i.e., the orthoborates $B(OR)_3$, $(RO)B(OR')_2$ and $(RO)B(OR')(OR'')$, and orthoborates of polyhydric alcohols (Greenwood et al., 1984). Organoboron compounds also include B-N compounds, because B-N is isoelectronic with C-C (Greenwood, 1973). Organoboron compounds are apparently important in biological systems and they are the result of interaction with OH or amine groups. As described below, organoboron complexes occur in plants and are produced in vitro with biomolecules isolated from animal tissues.

The most probable form of dietary boron after ingestion and subsequent hydrolysis (Greenwood, et al., 1984) is orthoboric acid (common name: boric acid) $B(OH)_3$. Boric acid accepts a hydroxyl ion (a Lewis acid) to form the tetrahedral anion $B(OH)_4^-$ (Reaction 1) (Greenwood, 1973):




At typical physiological boron concentrations (0.006 - ~9.0 $\mu\text{mol/L}$) in plants, animals, or humans, inorganic boron is essentially present only as the mononuclear species boric acid $B(OH)_3$ and as borate $B(OH)_4^-$ (Weser, 1967). Within the normal pH range of the gut and kidney, $B(OH)_3$ prevails as the dominant species (pH 1: ~100% $B(OH)_3$; pH 9.3: 50%; pH 11: ~0%) (Spivack et al., 1987).


2.2 Boron biochemistry

2.2.1 Boron esters


There is a vast number of mono or polyhydroxy compounds that contain one or more hydroxy groups with suitable positions for interaction with boron to form borate esters, specific organoboron complexes. For example, boric acid reacts with a suitable dihydroxy compound to form the corresponding boric acid monoester ("partial" esterification) (e.g., Structure 1) that retains the trigonal-planar configuration and no charge. Borate may react with a suitable dihydroxy compound to form the corresponding borate monoester ("partial" esterification; monocyclic) (Structure 2) with a tetrahedral configuration and a negative charge. A compound of similar configuration and charge is also formed when a boric acid monoester forms a complex with an available hydroxyl group. These two types of boromonoesters can react with another dihydroxy compound to give a corresponding spiro-cyclic borodiester ("complete" esterification) that is a chelate complex with a tetrahedral configuration and negative charge (Structure 3) (Van Duin et al., 1984).



(Structure 1)




(Structure 2)



(Structure 3)

2.2.2 Boron-Containing Biomolecules


Since discovery of the first boron biomolecule in 1967 (Hutter et al., 1967), several other similar biomolecules are now well-characterized (Loomis et al., 1992; O'Neill et al., 1996; Schummer et al., 1994). Most recently described was a cell-to-cell communication signal that requires boron (Chen et al., 2002). This cell signalling biomolecule, autoinducer-II (AI-II), is produced by a bacterium and binds to the primary receptor to form a furanosyl borate diester complex (Structure 4).



(Structure 4)

The discovery of all currently recognized boron-dependent biomolecules was achieved because the bound boron formed four coordinate covalent bonds with the ligand, creating a thermodynamically stable complex that is almost undissociable in water (Gerrard, 1961; Thellier et al., 1979).

Ribose derivatives have a strong affinity for boron (Loomis, et al., 1992) and nucleotide-boron adduction has been demonstrated for a number of ribose-containing nucleotides and cofactors, all with different affinities for boron (Structure 5) (Ralston et al., 2001). S-adenosylmethionine (SAM), the predominant methyl donor in biological methylations and a versatile cofactor in a variety of physiological processes, has the highest known affinity for boron of all animal and human biocompounds examined (Ralston, et al., 2001). Next in rank are members of the diadenosine phosphate (Ap_nA) family of biomolecules Ap_3A , Ap_4A , Ap_5A , and Ap_6A , signal nucleotides present in all cells with active protein synthesis (McLennan, 1992).



(Structure 5)

The physiological consequences of boron binding with SAM or the Ap_nA molecular species remain unexplored but probably are not trivial. NAD^+ (Structure 5) follows Ap_3A in rank for boron affinity (Ralston, et al., 2001) and oxidoreductase enzymes that require pyridine (e.g., NAD^+ or $NADP$) or flavin (e.g., FAD) nucleotides are well known to be competitively inhibited by borate or its derivatives (Kim et al., 2003; Smith et al., 1976). Reversible enzymatic inhibition as an essential role for an element is unusual. However, there is irrefutable evidence that boron serves to inhibit or dampen several metabolic pathways in plants (Lovatt et al., 1984; Shkol'nik et al., 1975).

Based on the structure of known boron-containing biomolecules, this author predicts that several biomolecules waiting discovery are derived from ribose and serve as signaling molecules that interact with the cell surface. Because known boron biomolecules are comprised of mirror or near-mirror halves stabilized by a single boron atom, it is predicted that undiscovered biomolecules have similar mirror construction.

3 BORON CONCENTRATIONS IN HEALTHY TISSUES

Boron is present at comparable concentrations in healthy tissues of different animals. Similar concentrations of boron ($\mu\text{g/mL}$) were reported in the plasma of humans (0.017-0.191) (Ferrando et al., 1993; Hunt et al., 1997; Iyengar et al., 1990; Mauras et al., 1986; Usuda et al., 1997; Wallace et al., 2002; Ward, 1993), rats (0.038-0.039) (Vaziri et al., 2001), chicks (0.047-0.152) (Hunt, 1996), cows (0.052-0.153) (Small et al., 1997), lambs (0.163) (Hunt, 1996), pigs (0.126) (Hunt, 1996), and horses (0.227) (Hunt, 1996). Liver boron concentrations ($\mu\text{g/g}$; dry weight) are similar in humans (1.1-5.4) (Shuler et al., 1990; Ward, 1987), rats (0.51) (Bai et al., 1996), chicks (1.01-4.4) (Rossi et al., 1993), and cows (3.3) (Ward, 1987); for brain tissue ($\mu\text{g/g}$; dry weight), similar in humans (0.87) (Shuler, et al., 1990), rats (0.64) (Bai, et al., 1996), and chicks (1.01-1.05); for

bone tissue ($\mu\text{g/g}$; dry weight), similar in humans (1.6) (Ward, 1993), rats (1.3) (Bai, et al., 1996), chicks (0.59-0.64) (Wilson et al., 1996), and mule deer (1.7) (Stelter, 1980).

4 ENVIRONMENTAL BORON

4.1 Dietary intakes of boron

Boron is ubiquitous in the environment and, on a molar basis, adult Americans consume more boron than several essential trace elements, e.g., copper, manganese, and molybdenum (Hunt et al., 2001). Boron consumption varies considerably among individuals and by sex-age group. For example, boron intake for infants aged 0 to 6 months is 0.75 ± 0.14 mg/d (mean \pm SE) (1st percentile, 0.03; 99th percentile, 6.40 mg/d); for males aged 51 to 70 years, 1.34 ± 0.02 mg/d (1st percentile, 0.39; 99th percentile, 3.34 mg/d); for lactating females, 1.39 ± 0.16 mg/d (1st percentile, 0.38; 99th percentile, 3.49 mg/d) (Food and Nutrition Board: Institute of Medicine, 2001). The range of dietary boron intakes within a sex-age group arises from a variety of factors. For example, compared with animal-based food products, plant-based products are much richer sources of dietary boron (Hunt, et al., 2001). Furthermore, most plant species within the subclass *Dicotyledoneae*, which includes fruits [i.e., raw pears: $2.27 \mu\text{g}$ ($0.21 \mu\text{mol}$) B/g], vegetables, tubers and legumes have much higher concentrations of boron than do species from the subclass *Monocotyledoneae*, especially gramineaceous species (the grasses) including rice [$0.09 \mu\text{g}$ ($0.008 \mu\text{mol}$) B/g], corn, barley, and wheat (Hunt, et al., 2001). For this reason, there are many diets otherwise nutritionally adequate that provide only 0.36 mg B/2000 kcal and are prepared easily by excluding vegetables, tubers, nuts, and legumes (Hunt, et al., 1997).

4.2 Boron toxicity

Boron has a range of safe exposures and it produces toxicity in all tested biological organisms when excessive amounts are absorbed. Boron dose response experiments to determine embryo-larval malformations in the frog *X. laevis* have demonstrated the pattern of areas of survival, deficiency, optimization, toxicity, and lethality that are characteristic of an essential element (Fort, et al., 2002).

Gastrointestinal absorption of boron approaches 100% (Hunt, et al., 1997). Even so, boron has a low order of toxicity. For adults, the Tolerable Upper Intake Level (UL) for boron is 20 mg/d (Food and Nutrition Board: Institute of Medicine, 2001), i.e., 20-fold typical intakes. Potentially lethal doses are generally cited as 3 g (3,000 mg) to 6 g (6,000 mg) for infants and 15 g (15,000 mg) to 20 g (20,000 mg) of boric acid for adults (Litovitz et al., 1988). At pre-

sent, death from boron poisoning is exceptionally rare probably because of the emphasis placed on maintaining electrolytic balance and supporting kidney function during the worst part of the poisoning incident.

5 BORON HOMEOSTATIC MECHANISMS

5.1 Boron gradient concentrations

There are several lines of evidence that suggest that boron may be regulated in humans. That boron contents in human milks were similar and stable throughout lactation of full term infants in two cohorts of women living in either Houston, TX, (Hunt et al., 2005) or St. John's, Newfoundland (Figure 1) (Hunt et al., 2004) has been interpreted as suggestive of regulatory mechanisms for the element, which remain undefined.

Substantial increases in dietary boron in elderly women [0.3 mg/d (1st percentile of typical intake) to 3.0 mg/d (~99th percentile of typical intake)] resulted in only small changes in blood boron values (Hunt, et al., 1997). Other investigators have also reported a remarkably narrow range of boron concentrations in whole blood from subjects with unknown dietary histories (Clarke et al., 1987).




Figure 1. Model and mean (\pm SE) concentrations of boron in breast milk from mothers of full-term (FT) and premature (PRT) infants; $n = 9$ per group over the 12 wk after birth. During the first 12 wk of lactation, prematurity affected the rate of change in concentrations ($P = 0.01$).

Work with higher mammals also suggests that boron may be regulated. In female rats, supplementation with high amounts of boron (9.25 mmol/L water) for 21 days, caused an increase in plasma boron concentrations but an undefined homeostatic mechanism concurrently eliminated any excess of boron from the liver and brain against their own concentration gradients (Magour et al., 1982). In yearling beef heifers, the percent of filtered boron reabsorbed by the kidneys decreased significantly with increased boron intake (Green et al., 1977). In gen-

eral, the findings to present suggest the presence of boron concentration against a gradient across mammalian cell membranes.

5.2 Boron transporters

The recent report (Ralston et al., 2004) that cultures of either RAW 264.7 cells or HL60 cells retain intracellular boron against a concentration gradient indicated the presence of intracellular boron binding species or the existence of boron specific transporters on the plasma membrane. Most likely, the repeated demonstration of the concentration of boron against a gradient indicates the existence of boron specific transporters. This line of evidence for the homeostatic control of boron is enhanced further by the discovery of a specific mammalian borate transporter, NaBC1, expressed in the basolateral membranes of epithelial cells (Park et al., 2004). The recent identification of the boron transporter, BOR1 (AtBor1), in the flowering plant *Arabidopsis thaliana* (Takano et al., 2005; Takano et al., 2002) and its mammalian homolog, BTR1, a newly discovered bicarbonate transporter superfamily member (Parker et al., 2001), provides further evidence for the homeostatic control of boron in humans and mammals.

6 BORON DEPLETION AND THE LIFE CYCLE

There are several lines of evidence that suggest that boron depletion prevents growth and completion of the life cycle in animal models. The original finding (Hunt et al., 1981) that boron deprivation can impair growth has been the basis for further research in several independent laboratories. Boron depletion can prevent growth and completion of the life cycle in the zebrafish. For example, in experiments with zebrafish, sperm from low-boron males successfully fertilized eggs from low-boron females (Rowe et al., 1999), but 92% of the embryos (compared to 37% of controls) died within 10 days. However, the low-boron embryos could be rescued from death if repleted with boron during the first hour after fertilization. Studies with the South African clawed frog, *Xenopus laevis* (Fort, et al., 2002), indicate that specimens fed a low-boron diet in a low-boron culture media produced a substantially higher number of necrotic eggs and fertilized embryos than frogs fed a boron-sufficient diet. By 96 hours of development, none of the larvae from boron-deficient adults maintained in low-boron culture media developed normally. Similar findings were reported for rats in which a low-boron diet (0.04 μ g/g) reduced the number of implantation sites compared to a diet supplemented with boron (Lanoue et al., 1998).

7 BORON DEPLETION AND PHYSIOLOGICAL/STRUCTURAL ABNORMALITIES

One of the criteria of essentiality, as defined by WHO, is a reduction in a physiologically important function when reduction in element exposure falls below a certain limit (Expert Consultation WHO/FAO/IAEA, 1996). Several lines of evidence indicate the boron deprivation appears to perturb physiological processes in frogs, zebrafish, chicks, rats, pigs, and humans.

7.1 Boron and calcium metabolism and bone growth and maintenance

Observations in human studies suggest that boron can influence calcium metabolism. For example, an increase in boron intake (0.36 to 3.23 mg/d) by postmenopausal women resulted in a 5% increase in urinary calcium excretion (Hunt, et al., 1997). Because increases in dietary calcium often result in increased urinary calcium excretion, this finding may reflect an increase in intestinal calcium absorption. Thus, it is important to determine whether the primary effect of boron on calcium metabolism is at the level of enteric absorption.

Bone calcification and metabolism respond to boron nutrition. Maturation of the growth plate was retarded during dietary boron deprivation in the chick (Hunt et al., 1994). Boron deprivation reduced bone strength in pigs (Armstrong et al., 2000) and rats (Nielsen, 2004) and induced abnormal limb development in frogs (Fort et al., 2000). In other studies, a boron supplement increased bone strength in nutritionally adequate chicks (Wilson et al., 1998) and adult rats (Chapin et al., 1998).

There is considerable evidence that dietary boron alleviates the signs of marginal vitamin D deficiency. Marginal vitamin D deficiency elevates plasma alkaline phosphatase concentrations, reduces body weight, and impairs bone structure. In the growing rachitic chick, dietary boron reduced elevated serum concentrations of alkaline phosphatase (Hunt, et al., 1981; Kurtoglu et al., 2001), improved body weight (Bai et al., 1996; Kurtoglu, et al., 2001), and substantially alleviated the perturbed histomorphometric indices of bone growth cartilage (Hunt, 1989; Hunt, et al., 1994).

7.2 Dietary boron and insulin metabolism

Pancreatic β -cells easily induced to secrete mass quantities of insulin are readily damaged, which eventually can cause them to stop functioning and result in diabetes mellitus (Reaven, 1999; Sprietsma et al., 1993). It is possible that dietary boron may reduce the amount of insulin needed to maintain glucose levels, thus limiting β -cell deterioration. For

example, as a dietary ingredient, boron decreased peak pancreatic in situ insulin release in chicks (Figure 2). In rats, dietary boron decreased plasma insulin concentrations but did not change glucose concentrations (Bakken et al., 2003). Work is underway to determine whether dietary boron can reduce β -cell "exhaustion" and deterioration.




Figure 2. Alteration in peak insulin secretion from isolated, perfused pancreata from 1-d-old cockerels fed a diet containing 0.29 (boron-low) or 1.65 mg B/kg and supplemented with cholecalciferol (Vit D) at 3.13 (inadequate) or 15.60 (adequate) $\mu\text{g/kg}$ for 26-37 d. Perfusion phases varied in the total amount of glucose (G: 5.5 or 38.0 mmol/L) and boron (B; 0.9 or 13.2 $\mu\text{mol/L}$) in the perfusate.

8 CONCLUSIONS

Boron is a natural constituent of the diet. Adult Americans consume slightly less than 1.0 mg per day on average and can easily increase that average by increasing consumption of fruits and vegetables. The evidence to date suggests that humans and at least some higher animals may use boron to support normal biological functions. The biochemical mechanisms responsible for these effects are poorly understood. However, the nature of boron biochemistry suggests specific lines of investigation. In particular, further characterization of the cell signalling molecules capable of complexing with boron should provide insights into the specific biochemical function(s) of boron in humans.

REFERENCES

- Armstrong, T.A., Spears, J.W., Crenshaw, T.D. and Nielsen, F.H. 2000. Boron supplementation of a semipurified diet for weanling pigs improves feed efficiency and bone strength characteristics and alters plasma lipid metabolites. *J. Nutr.* 130:2575-2581.
- Bai, Y. and Hunt, C.D. 1996. Absorption and distribution of boron in rats following a single oral administration of boron. *Proc. N. D. Acad. Sci.* 50:53.
- Bai, Y. and Hunt, C.D. 1996. Dietary boron enhances efficacy of cholecalciferol in broiler chicks. *J. Trace Elem. Exp. Med* 9:117-132.

- Bakken, N.A. and Hunt, C.D. 2003. Dietary boron decreases peak pancreatic in situ insulin release in chicks and plasma insulin concentrations in rats regardless of vitamin D or magnesium status. *J. Nutr.* 133:3516-3522.
- Carney, G.E. and Bowen, N.J. 2004. p24 proteins, intracellular trafficking, and behavior: *Drosophila melanogaster* provides insights and opportunities. *Biol. Cell* 96:271-8.
- Chapin, R.E., Ku, W.W., Kenney, M.A. and McCoy, H. 1998. The effects of dietary boric acid on bone strength in rats. *Biol. Trace Elem. Res.* 66:395-399.
- Chen, T.S.S., Chang, C.-j. and Floss, H.G. 1981. On the biosynthesis of boromycin. *J. Org. Chem.* 46:2661-2665.
- Chen, X., Schauder, S., Potier, N., Van Dorsselaer, A., Pelczar, I., Bassler, B.L. and Hughson, F.M. 2002. Structural identification of a bacterial quorum-sensing signal containing boron. *Nature* 415:545-549.
- Clarke, W.B., Koekebakker, M., Barr, R.D., Downing, R.G. and Fleming, R.F. 1987. Analysis of ultratrace lithium and boron by neutron activation and mass-spectrometric measurement of ^3He and ^4He . *Int. J. Rad. Appl. Instrum.* 38:735-743.
- Expert Consultation WHO/FAO/IAEA 1996. Trace elements in human nutrition and health. Geneva:WHO.
- Ferrando, A.A., Green, N.R., Barnes, K.W. and Woodward, B. 1993. Microwave digestion preparation and ICP determination of boron in human plasma. *Biol. Trace Elem. Res.* 37:17-25.
- Food and Nutrition Board: Institute of Medicine 2001. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. Washington, D.C.:National Academy Press.
- Fort, D.J., Rogers, R.L., McLaughlin, D.W., Sellers, C.M. and Schlekut, C.L. 2002. Impact of boron deficiency on *Xenopus laevis*: a summary of biological effects and potential biochemical roles. *Biol. Trace Elem. Res.* 90:117-42.
- Fort, D.J., Stover, E.L., Rogers, R.L., Copley, H.F., Morgan, L.A. and Foster, E.R. 2000. Chronic boron or copper deficiency induces limb teratogenesis in *Xenopus*. *Biol. Trace Elem. Res.* 77:173-87.
- Frieden, E. 1984. A survey of the essential biochemical elements. In: *Biochemistry of the essential ultratrace elements*, (Frieden, E, eds). New York:Plenum Press.
- Gerrard, W. 1961. The Organic Chemistry of Boron. London:Academic Press.
- Green, G.H. and Weeth, H.J. 1977. Responses of heifers ingesting boron in water. *J. Anim. Sci.* 46:812-818.
- Greenwood, N.N. 1973. Boron. In: *Comprehensive Inorganic Chemistry*, (Bailar, J. J., Emeléus, H., Nyholm, R and Trotman-Dickenson, A, eds). Oxford:Pergamon Press Ltd.
- Greenwood, N.N. and Earnshaw, A. 1984. Chemistry of the Elements. Oxford, Great Britain:Pergamon Press.
- Hunt, C.D. 1989. Dietary boron modified the effects of magnesium and molybdenum on mineral metabolism in the cholecalciferol-deficient chick. *Biol. Trace Elem. Res.* 22:201-220.
- Hunt, C.D. 1996. Dietary boron deficiency and supplementation. In: *Trace Elements in Laboratory Rodents*, (Watson, R R, eds). Boca Raton, FL:CRC Press.
- Hunt, C.D., Butte, N.F. and Johnson, L.K. 2005. Boron concentrations in milk from mothers of exclusively breast-fed healthy full-term infants are stable during the first four months of lactation. *J. Nutr.* 135:2383-2386.
- Hunt, C.D., Friel, J.K. and Johnson, L.K. 2004. Boron concentrations in milk from mothers of full-term and premature infants. *Am. J. Clin. Nutr.* 80:1327-33.
- Hunt, C.D., Herbel, J.L. and Idso, J.P. 1994. Dietary boron modifies the effects of vitamin D₃ nutriture on indices of energy substrate utilization and mineral metabolism in the chick. *J. Bone Miner. Res.* 9:171-181.
- Hunt, C.D., Herbel, J.L. and Nielsen, F.H. 1997. Metabolic response of postmenopausal women to supplemental dietary boron and aluminum during usual and low magnesium intake: boron, calcium, and magnesium absorption and retention and blood mineral concentrations. *Am. J. Clin. Nutr.* 65:803-813.
- Hunt, C.D. and Meacham, S.L. 2001. Aluminum, boron, calcium, copper, iron, magnesium, manganese, molybdenum, phosphorus, potassium, sodium, and zinc: concentrations in common Western foods and estimated daily intakes by infants, toddlers, and male and female adolescents, adults, and seniors in the United States. *J. Am. Diet. Assoc.* 101:1058-1060; Table, http://www.eatright.com/images/journal/0901/hunt_tables.pdf.
- Hunt, C.D. and Nielsen, F.H. 1981. Interaction between boron and cholecalciferol in the chick. In: *Trace Element Metabolism in Man and Animals-4*, (Gawthorne, J and White, C, eds). Canberra:Australian Academy of Science.
- Hutter, R., Keller-Schierlein, W., Knusel, F., Prelog, V., Rodgers, G.C., Jr., Suter, P., Vogel, G., Voser, W. and Zahner, H. 1967. Stoffwechselprodukte von Mikroorganismen. Boromycin (The metabolic products of microorganisms. Boromycin). *Helv. Chim. Acta* 50:1533-9.
- Iyengar, G.V., Clarke, W.B. and Downing, R.G. 1990. Determination of boron and lithium in diverse biological matrices using neutron activation-mass spectrometry (NA-MS). *Fres. J. Anal. Chem.* 338:562-566.
- Kim, D.H., Marbois, B.N., Faull, K.F. and Eckhart, C.D. 2003. Esterification of borate with NAD⁺ and NADH as studied by electrospray ionization mass spectrometry and ¹¹B NMR spectroscopy. *J. Mass Spectrom.* 38:632-40.
- Kurtoglu, V., Kurtoglu, F. and Coskun, B. 2001. Effects of boron supplementation of adequate and inadequate vitamin D₃- containing diet on performance and serum biochemical characters of broiler chickens. *Res. Vet. Sci.* 71:183-187.
- Lanoue, L., Taubeneck, M.W., Muniz, J., Hanna, L.A., Strong, P.L., Murray, F.J., Nielsen, F.H., Hunt, C.D. and Keen, C.L. 1998. Assessing the effects of low boron diets on embryonic and fetal development in rodents using in vitro and in vivo model systems. *Biol. Trace Elem. Res.* 66:271-298.
- Litovitz, T.L., Klein-Schwartz, W., Oderda, G.M. and Schmitz, B.F. 1988. Clinical manifestations of toxicity in a series of 784 boric acid ingestions. *Am. J. Emerg. Med.* 6:209-213.
- Loomis, W.D. and Durst, R.W. 1992. Chemistry and biology of boron. *BioFactors* 3:229-39.

- Lovatt, C.J. 1985. Evolution of xylem resulted in a requirement for boron in the apical meristems of vascular plants. *New Phytol.* 99:509-522.
- Lovatt, C.J. and Dugger, W.M. 1984. Boron. In: *Biochemistry of the Essential Ultratrace Elements*, (Frieden, E, eds). New York:Plenum Press.
- Magour, S., Schramel, P., Ovcar, J. and Mäser, H. 1982. Uptake and distribution of boron in rats: interaction with ethanol and hexobarbital in the brain. *Arch. Environm. Contam. Toxicol.* 11:521-525.
- Mauras, Y., ANg, K.S., Simon, P., Tessier, B., Cartier, F. and Allain, P. 1986. Increase in blood plasma levels of boron and strontium in hemodialyzed patients. *Clin. Chim. Acta* 156:315-320.
- McLennan, A.G. 1992. Dinucleoside phosphates--an introduction. In: *ApA and other dinucleoside polyphosphates*, (McLennan, A G, eds). Boca Raton:CRC Press.
- Mertz, W. 1970. Some aspects of nutritional trace element research. *Fed. Proc.* 29:1482-1488.
- Nielsen, F.H. 2004. Dietary fat composition modifies the effect of boron on bone characteristics and plasma lipids in rats. *Biofactors* 20:161-71.
- O'Neill, M.A., Warrenfeltz, D., Kates, K., Pellerin, P., Doco, T., Darvill, A.G. and Albersheim, P. 1996. Rhamnogalacturonan-II, a pectic polysaccharide in the walls of growing plant cells, forms a dimer that is covalently cross-linked by a borate ester. *J. Biol. Chem.* 271:22923-22930.
- Park, M., Li, Q., Shcheynikov, N., Zeng, W. and Muallem, S. 2004. NaBC1 is a ubiquitous electrogenic Na(+)-coupled borate transporter essential for cellular boron homeostasis and cell growth and proliferation. *Mol Cell* 16:331-41.
- Parker, M.D., Ourmozdi, E.P. and Tanner, M.J. 2001. Human BTR1, a new bicarbonate transporter superfamily member and human AE4 from kidney. *Biochem Biophys Res Commun* 282:1103-9.
- Ralston, N.V.C. and Hunt, C.D. 2001. Diadenosine phosphates and S-adenosylmethionine: novel boron binding biomolecules detected by capillary electrophoresis. *Biochim. Biophys. Acta* 1527:20-30.
- Ralston, N.V.C. and Hunt, C.D. 2004. Transmembrane partitioning of boron and other elements in RAW 264.7 and HL60 cell cultures. *Biol. Trace Elem. Res.* 98:181-192.
- Reaven, G.M. 1999. Insulin resistance: a chicken that has come to roost. *Ann N Y Acad Sci* 892:45-57.
- Rossi, A.F., Miles, R.D., Damron, B.L. and Flunker, L.K. 1993. Effects of dietary boron supplementation on broilers. *Poul. Sci.* 72:2124-2130.
- Rowe, G. 1999. Surveying boron and naturally soft groundwater with a computer database. *J. Environ. Health* 62:14-18.
- Rowe, R.I. and Eckhart, C.D. 1999. Boron is required for zebrafish embryogenesis. *J. Exp. Biol.* 202:1649-1654.
- Schummer, D., Irschik, H., Reichenbach, H. and Höfle, G. 1994. Antibiotics from gliding bacteria, LVII. Tartrolons: new boron-containing macrodiolides from *Sorangium cellulosum*. *Liebigs Ann. Chem.* 1994:283-289.
- Shkolnik, M.Y. and Il'inskaya, N.L. 1975. Effect of a boron deficiency on activity of glucose-6-phosphate dehydrogenase in plants with different boron requirements. *Fiziol. Rastenii* 22:801-805.
- Shuler, T.R., Pootrakul, P., Yarnsukon, P. and Nielsen, F.H. 1990. Effect of thalassaemia/immunoglobulin E disease on macro, trace and ultratrace element concentrations in human tissues. *J. Trace Elem. Exp. Med.* 3:31-43.
- Small, J.A., Charmley, E., Rodd, A.V. and Fredeen, A.H. 1997. Serum mineral concentrations in relation to estrus and conception in beef heifers and cows fed conserved forage. *Can. J. Anim. Sci.* 77:55-62.
- Smith, K.W. and Johnson, S.L. 1976. Borate inhibition of yeast alcohol dehydrogenase. *Biochem.* 15:560-565.
- Spivack, A.J. and Edmond, J.M. 1987. Boron isotope exchange between seawater and the oceanic crust. *Geochim. Cosmochim. Acta* 51:1033-1043.
- Sprietsma, J. and Schuitemaker, G. 1993. Diabetes can be prevented by reducing insulin production. *Med. Hypotheses* 42:15-23.
- Stelter, L.H. 1980. Baseline levels of selected trace elements in Colorado oil shale region animals. *J. Wildlife Dis.* 16:175-182.
- Takano, J., Miwa, K., Yuan, L., von Wiren, N. and Fujiwara, T. 2005. Endocytosis and degradation of BOR1, a boron transporter of Arabidopsis thaliana, regulated by boron availability. *Proc Natl Acad Sci U S A* 102:12276-81.
- Takano, J., Noguchi, K., Yasumori, M., Kobayashi, M., Gajdos, Z., Miwa, K., Hayashi, H., Yoneyama, T. and Fujiwara, T. 2002. Arabidopsis boron transporter for xylem loading. *Nature* 420:337-40.
- Thellier, M., Duval, Y. and Demarty, M. 1979. Borate exchanges of *Lemna minor* L. as studied with the help of the enriched stable isotopes and of a (n, alpha) nuclear reaction. *Plant Physiol.* 63:283-288.
- Underwood, E.J. and Mertz, W. 1987. Introduction. In: *Trace Elements in Human and Animal Nutrition--Fifth Edition*, (Mertz, W, eds). San Diego:Academic Press.
- Usuda, K., Kono, K. and Yoshida, Y. 1997. Serum boron concentration from inhabitants of an urban area in Japan. *Biol. Trace Ele. Res.* 56:167-178.
- Van Duin, M., Peters, J.A., Kieboom, A.P.G. and Van Bakkum, H. 1984. Studies on borate esters I. The pH dependence of the stability of esters of boric acid and borate in aqueous medium as studied by ¹¹B NMR. *Tetrahedron* 40:2901-2911.
- Vaziri, N.D., Oveisi, F., Culver, B.D., Pahl, M.V., Andersen, M.E., Strong, P.L. and Murray, F.J. 2001. The effect of pregnancy on renal clearance of boron in rats given boric acid orally. *Toxicol Sci* 60:257-63.
- Wallace, J.M., Hannon-Fletcher, M.P., Robson, P.J., Gilmore, W.S., Hubbard, S.A. and Strain, J.J. 2002. Boron supplementation and activated factor VII in healthy men. *Eur J Clin Nutr* 56:1102-7.
- Ward, N.I. 1987. The determination of boron in biological materials by neutron irradiation and prompt gamma-ray spectrometry. *J. Rad. Nuc. Chem.* 110:633-639.
- Ward, N.I. 1993. Boron levels in human tissues and fluids. In: *Trace Element Metabolism in Man and Animals-8*, (Anke, M, Meissner, D and Mills, C, eds). Gersdorf, Germany:Verlag Media Touristik.

- Warington, K. 1926. The changes induced in the anatomical structure of *Vicia Faba* by the absence of boron from the nutrient solution. *Ann. Bot.* 40:27-42.
- Weser, U. 1967. Chemistry and structure of some borate polyol compounds of biochemical interest. In: *Structure and Bonding*, (Jorgensen, C, Neilands, J, Nyholm, R, Reinen, D and Williams, R, eds). New York:Springer-Verlag.
- Wilson, J.H. and Ruzler, P.L. 1996. Effects of dietary boron supplementation on laying hens. *Br Poult Sci* 37:723-9.
- Wilson, J.H. and Ruzler, P.L. 1998. Long term effects of boron on layer bone strength and production parameters. *Br Poult Sci* 39:11-5.

III. ULUSLARARASI BOR SEMPOZYUMU BİLDİRİLER KİTABI

PROCEEDINGS OF 3rd INTERNATIONAL BORON SYMPOSIUM

02-04 Kasım/November
2006 Ankara, TÜRKİYE

Edi.t.örler/Editors

Tevfik GÜYAGÜLER
Sinem KARAKAŞ
Zeynep ÖZDEMİR
Ahmet KARAKAŞ
Ayşe BÜKÜLMEZ
Oğuz SÖNMEZER



TMMOB Maden Mühendisleri Odası
The Chamber of
Mining Engineers of Turkey



Eti Maden İşletmeleri
Genel Müdürlüğü
Eti Mine Works General Management

BORON

Ulusal Bor Araştırma Enstitüsü
National Boron
Research Institute

Tüm hakları saklıdır © 2006

TMMOB Maden Mühendisleri Odası'nın yazılı izni olmaksızın bu kitap ya da bu kitabın bir kısmı herhangi bir biçimde yayınlanamaz.

All right reserved © 2006

No parts of this book may be reproduced in any form or any means, without written permission of the Chamber of Mining Engineers of Turkey.

Baskı / Published by Kozan Ofset Matbaacılık

Büyük Sanayi 1. Cadde No: 95/11 İskitler / ANKARA

Tel : +90 (312) 384 2003 Fax : +90 (312) 341 2860

ISBN 9944-89-182-7

TMMOB – Maden Mühendisleri Odası / *The Chamber of Mining Engineers of Turkey*

Selânik Cad. No: 19/4 06650 – Kızılay, Ankara, Turkey

Tel: +90 (312) 425 10 80 Fax : +90 (312) 417 59 90

Web : www.maden.org.tr E-mail: maden@maden.org.tr